arXiv:gr-qc/0512055v1 9 Dec 2005

Analytic expressions for gravitational inner multipole moments of elementary solids and for the force between two rectangular solids

E. G. Adelberger, Nathan A. Collins‡ **and C.D. Hoyle**

Department of Physics, University of Washington, Seattle, Washington 98195-1560

E-mail: `hoyle@npl.washington.edu`

Abstract. We give analytic expressions for the gravitational inner spherical multipole moments, q_{lm} with $l \leq 5$, for eleven elementary solid shapes. These moments, in conjunction with their known rotational and translational properties, can be used to calculate precisely the moments of complex objects that may be assembled from the elementary shapes. We also give an analytic expression for the gravitational force between two rectangular solids at arbitrary separations. These expressions are useful for computing the gravitational properties of complex instruments, such as those used in equivalence principle tests, and in the gravitational balancing of drag-free spacecraft.

PACS numbers: 04.20.Cv,04.80.-y

Submitted to: *Class. Quantum Grav.*

---

‡ Present address: Encina Hall West, Room 100, Stanford University, Stanford, CA, 94305-6044

1. Introduction

As the precision of gravitational experiments increases it is increasingly necessary to compute with high accuracy the gravitational fields of complicated objects. This paper provides analytic results that have proved useful for this task.

Multipole expansions provide a powerful formalism useful in many physical applications. One such application occurs in equivalence principle tests [1, 2]where gravity gradients set a fundamental limit to the precision of the results. Spurious effects from gravity gradients can be minimized in a systematic way by nulling the low-order multipole moments of the differential accelerometer (typically a torsion pendulum) and the attractor. In the case of torsion balances, where differential acceleration of the test bodies produces a torque on the pendulum, expansions in spherical multipoles are particularly useful. Evaluating the multipole moments of complicated objects presents a challenge. The moments can be found by decomposing the objects into elementary 3-dimensional shapes whose moments can be calculated exactly. Because the moments have definite rotational and translational properties[3], a catalog of moments of the elementary solids evaluated in any suitable coordinate system allows one to "assemble" realistic objects and compute their multipole moments with high precision. The formalism of Ref. [3], together with the moments cataloged here, provides a convenient way to understand the gravitational response of torsion balances, including the effects of imperfections in the instruments themselves. This paper gives analytic expressions for the inner multipole moments of 11 elementary solid objects that have proved useful in modeling the gravitational response of actual devices.

Another application of analytic results is found in the gravitational compensation of drag-free spacecraft. For example, the LISA (Laser Interferometer Space Antenna) gravitational wave detector will use three such spacecraft, each containing two inertial bodies that form the end mirrors of interferometer arms. Stringent requirements on the test-body DC acceleration and force noise require accurate modeling of gravitational forces, torques, and gradients produced by the surrounding spacecraft enclosure [4]. Since much of the disturbing mass distribution is in close proximity to the inertial bodies, the multipole formalism is not valid for this task. The analytic expression for the force between two rectangular solids, derived here, provides an accurate tool for assessing these gravitational disturbances.

2. Inner moments of the elementary solids

We present analytic expressions for the gravitational (or Coulomb) inner moments, q_{lm},

$$q_{lm} = \rho \int r^l Y^*_{lm}(\hat{r}) d^3r \ ,$$

of the following elementary shapes: sphere, rectangular solid, cylindrical annulus, cone, triangular slab, trapezoidal slab, cylindrical section, right tetrahedron, hole in a cylinder, hole in a plate, and pyramid. All moments are given with respect to the origin, and are proportional to the mass density, ρ. We list all non-vanishing moments with $l \leq 5$; the unlisted $l \leq 5$ moments vanish identically.

2.1. Sphere

The sphere of radius r is centered on the origin.

$$q_{00} = \frac{2}{3}\rho\sqrt{\pi}r^3$$

2.2. Rectangular box

The box is centered on the origin and is specified by the positive quantities x, y, and z–its dimensions along $\hat{x}$, $\hat{y}$, and $\hat{z}$.

$$q_{00} = \frac{\rho xyz}{2\sqrt{\pi}}$$

$$q_{20} = -q_{00}\frac{1}{24}\sqrt{5}\left(x^2+y^2-2z^2\right)$$

$$q_{22} = q_{00}\frac{1}{8}\sqrt{\frac{5}{6}}\left(x^2-y^2\right)$$

$$q_{40} = q_{00}\frac{1}{640}\left[9x^4+10x^2y^2+9y^4-40\left(x^2+y^2\right)z^2+24z^4\right]$$

$$q_{42} = -q_{00}\frac{1}{64\sqrt{10}}\left(x^2-y^2\right)\left[3\left(x^2+y^2\right)-10z^2\right]$$

$$q_{44} = q_{00}\frac{1}{128}\sqrt{\frac{7}{10}}\left(3x^4-10x^2y^2+3y^4\right)$$

2.3. Cylindrical annulus

The cylindrical annulus has its axis of symmetry along $\hat{z}$ and is symmetric under reflections through the $\hat{x}$-$\hat{y}$ plane. It is specified by inner and outer radii, r_1 and r_2, and height $h > 0$. It need not be a complete annulus, but spans azimuthal angles between ϕ_1 and ϕ_2, which are measured with respect to $\hat{x}$. We define $\Delta_\Phi \equiv \phi_2 - \phi_1$, $\phi \equiv (\phi_1 + \phi_2)/2$, $\Delta_R \equiv r_2 - r_1$, and $r \equiv (r_1 + r_2)/2$.

$$q_{00} = \frac{\rho}{2\sqrt{\pi}}hr\Delta_\Phi\Delta_R$$

$$q_{11} = -\frac{\rho}{4\sqrt{6\pi}}e^{-i\phi}h\Delta_R\left(\Delta_R^2+12r^2\right)\sin\frac{\Delta_\Phi}{2}$$

$$q_{20} = -\frac{\rho}{48}\sqrt{\frac{5}{\pi}}hr\Delta_\Phi\Delta_R\left(3\Delta_R^2-2h^2+12r^2\right)$$

$$q_{22} = \frac{\rho}{16}\sqrt{\frac{15}{2\pi}}e^{-2i\phi}hr\Delta_R\left(\Delta_R^2+4r^2\right)\sin\Delta_\Phi$$

$$q_{31} = \frac{\rho}{960}\sqrt{\frac{7}{3\pi}}e^{-i\phi}h\Delta_R\Big[9\Delta_R^4-20\Delta_R^2\left(h^2-18r^2\right)-240r^2\left(h^2-3r^2\right)\Big]\sin\frac{\Delta_\Phi}{2}$$

$$q_{33} = -\frac{\rho}{192}\sqrt{\frac{7}{5\pi}}e^{-3i\phi}h\Delta_R\left(\Delta_R^4+40\Delta_R^2r^2+80r^4\right)\sin\frac{3\Delta_\Phi}{2}$$

$$q_{40} = \frac{3\rho}{1280\sqrt{\pi}}hr\Delta_\Phi\Delta_R\Big[15\Delta_R^4-40\Delta_R^2\left(h^2-5r^2\right)+8\left(h^4-20h^2r^2+30r^4\right)\Big]$$

$$q_{42} = -\frac{\rho}{128}\sqrt{\frac{5}{2\pi}}e^{-2i\phi}hr\Delta_R\Big[3\Delta_R^4-24r^2\left(h^2-2r^2\right)+\Delta_R^2\left(-6h^2+40r^2\right)\Big]\sin\Delta_\Phi$$

$$q_{44} = \frac{\rho}{512}\sqrt{\frac{35}{2\pi}}e^{-4i\phi}hr\Delta_R\big(3\Delta_R^4 + 40\Delta_R^2 r^2 + 48r^4\big)\sin 2\Delta_\Phi$$

$$\begin{aligned} q_{51} = -\frac{\rho}{3584}\sqrt{\frac{11}{30\pi}}e^{-i\phi}h\Delta_R\Big[&15\Delta_R^6 - 84\Delta_R^4\left(h^2 - 15r^2\right) + 672r^2\left(h^4 - 10h^2r^2 + 10r^4\right) \\ &+56\Delta_R^2\left(h^4 - 60h^2r^2 + 150r^4\right)\Big]\sin\frac{\Delta_\Phi}{2} \end{aligned}$$

$$\begin{aligned} q_{53} = \frac{\rho}{9216}\sqrt{\frac{11}{35\pi}}e^{-3i\phi}h\Delta_R\Big[&15\Delta_R^6 - 56\Delta_R^4 h^2 + 140\Delta_R^2\left(9\Delta_R^2 - 16h^2\right)r^2 \\ &+560\left(15\Delta_R^2 - 8h^2\right)r^4 + 6720r^6\Big]\sin\frac{3\Delta_\Phi}{2} \end{aligned}$$

$$q_{55} = -\frac{3\rho}{5120}\sqrt{\frac{11}{7\pi}}e^{-5i\phi}h\Delta_R\Big(\Delta_R^6 + 84\Delta_R^4 r^2 + 560\Delta_R^2 r^4 + 448r^6\Big)\sin\frac{5\Delta_\Phi}{2}$$

2.4. Cone

The (truncated) cone is symmetrical about $\hat{z}$, and is specified by upper and lower radii, r_1 and r_2, and height $h > 0$. The cone extends a distance $h/2$ above and below the $\hat{x}$-$\hat{y}$ plane. Complete cones have vanishing values of r_1 or r_2.

$$q_{00} = \frac{\rho}{6}\sqrt{\pi}h\left(r_1^2 + r_1r_2 + r_2^2\right)$$

$$q_{10} = \frac{\rho}{8}\sqrt{\frac{\pi}{3}}h^2\left(r_1^2 + 2r_1r_2 + 3r_2^2\right)$$

$$q_{20} = \frac{\rho}{24}\sqrt{\frac{\pi}{5}}h\Big[2h^2\left(r_1^2 + 3r_1r_2 + 6r_2^2\right) - 3\big(r_1^4 + r_1^3r_2 + r_1^2r_2^2 + r_1r_2^3 + r_2^4\big)\Big]$$

$$q_{30} = \frac{\rho}{240}\sqrt{7\pi}h^2\Big[2h^2\left(r_1^2 + 4r_1r_2 + 10r_2^2\right) - 3\big(r_1^4 + 2r_1^3r_2 + 3r_1^2r_2^2 + 4r_1r_2^3 + 5r_2^4\big)\Big]$$

$$\begin{aligned} q_{40} = \frac{\rho}{560}\sqrt{\pi}h\Big[&8h^4\left(r_1^2 + 5r_1r_2 + 15r_2^2\right) - 12h^2\left(r_1^4 + 3r_1^3r_2 + 6r_1^2r_2^2 + 10r_1r_2^3 + 15r_2^4\right) \\ &+15\left(r_1^6 + r_1^5r_2 + r_1^4r_2^2 + r_1^3r_2^3 + r_1^2r_2^4 + r_1r_2^5 + r_2^6\right)\Big] \end{aligned}$$

$$\begin{aligned} q_{50} = \frac{\rho}{2688}\sqrt{11\pi}h^2\Big[&8h^4\left(r_1^2 + 6r_1r_2 + 21r_2^2\right) - 12h^2\left(r_1^4 + 4r_1^3r_2 + 10r_1^2r_2^2 + 20r_1r_2^3 + 35r_2^4\right) \\ &+15\big(r_1^6 + 2r_1^5r_2 + 3r_1^4r_2^2 + 4r_1^3r_2^3 + 5r_1^2r_2^4 + 6r_1r_2^5 + 7r_2^6\big)\Big] \end{aligned}$$

2.5. Triangular slab

This shape has reflection symmetry about the $\hat{x}$-$\hat{y}$ plane and thickness $h > 0$. The triangular faces have vertices at $(x, y) = (0, 0)$, (d, y_1) and (d, y_2). The restriction that one side be parallel to $\hat{y}$ is easily overcome using the rotational properties of the moments [3].

$$q_{00} = \frac{\rho}{4\sqrt{\pi}}h|d\left(y_1 - y_2\right)|$$

$$q_{11} = q_{00}\left[-2d + i\left(y_1 + y_2\right)\right]$$

$$q_{20} = -q_{00}\frac{\sqrt{5}}{12}\left(3d^2 - h^2 + y_1^2 + y_1y_2 + y_2^2\right)$$

$$q_{22} = q_{00}\frac{1}{4}\sqrt{\frac{5}{6}}\Big[3d^2 - y_1^2 - y_1y_2 - y_2^2 - 3id\left(y_1 + y_2\right)\Big]$$

$$\begin{aligned}q_{31} = q_{00}\frac{1}{120}\sqrt{\frac{7}{3}}\Big\{&36d^3 - 18id^2\left(y_1+y_2\right) + i\left(y_1+y_2\right)\left[10h^2 - 9\left(y_1^2+y_2^2\right)\right]\\ &-4d\left[5h^2 - 3\left(y_1^2+y_1y_2+y_2^2\right)\right]\Big\}\end{aligned}$$

$$q_{33} = q_{00}\frac{1}{8}\sqrt{\frac{7}{5}}\left[(1+i)\,d + y_1 - iy_2\right] \times \left[(1+i)\,d - iy_1 + y_2\right]\left[2id + \left(y_1+y_2\right)\right]$$

$$\begin{aligned}q_{40} = q_{00}\frac{1}{80}\Big\{&30d^4 + 3h^4 - 10h^2\left(y_1^2+y_1y_2+y_2^2\right) + 6\left(y_1^4+y_1^3y_2+y_1^2y_2^2+y_1y_2^3+y_2^4\right)\\ &+10d^2\left[-3h^2 + 2\left(y_1^2+y_1y_2+y_2^2\right)\right]\Big\}\end{aligned}$$

$$\begin{aligned}q_{42} = -q_{00}\frac{1}{16\sqrt{10}}\Big\{&20d^4 - 15d^2h^2 - 20id^3\left(y_1+y_2\right) + 5h^2\left(y_1^2+y_1y_2+y_2^2\right)\\ &-4\left(y_1^4+y_1^3y_2+y_1^2y_2^2+y_1y_2^3+y_2^4\right) + 5id\left(y_1+y_2\right)\left[3h^2 - 2\left(y_1^2+y_2^2\right)\right]\Big\}\end{aligned}$$

$$\begin{aligned}q_{44} = q_{00}\frac{1}{8}\sqrt{\frac{7}{10}}\Big[&5d^4 + y_1^4 + y_1^3y_2 + y_1^2y_2^2 + y_1y_2^3 + y_2^4 - 10id^3\left(y_1+y_2\right)\\ &+5id\left(y_1+y_2\right)\left(y_1^2+y_2^2\right) - 10d^2\left(y_1^2+y_1y_2+y_2^2\right)\Big]\end{aligned}$$

$$\begin{aligned}q_{51} = -q_{00}\frac{1}{112}\sqrt{\frac{11}{30}}\Big\{&60d^5 - 30id^4(y_1+y_2) + 6id^2(y_1+y_2)\left[7h^2 - 5(y_1^2+y_2^2)\right]\\ &-4d^3\left[21h^2 - 10\left(y_1^2+y_1y_2+y_2^2\right)\right] + 2d\big[7h^4 - 14h^2\left(y_1^2+y_1y_2+y_2^2\right)\\ &+6\left(y_1^4+y_1^3y_2+y_1^2y_2^2+y_1y_2^3+y_2^4\right)\big]\\ &-i\big[7h^4(y_1+y_2) - 21h^2(y_1^3+y_1^2y_2+y_1y_2^2+y_2^3)\\ &+10\left(y_1^5+y_1^4y_2+y_1^3y_2^2+y_1^2y_2^3+y_1y_2^4+y_2^5\right)\big]\Big\}\end{aligned}$$

$$\begin{aligned}q_{53} = q_{00}\frac{1}{48}\sqrt{\frac{11}{35}}\Big\{&30d^5 - 45id^4\left(y_1+y_2\right) + 3id^2\left(y_1+y_2\right)\left[14h^2 - 5\left(y_1^2+y_2^2\right)\right]\\ &-4d^3\left[7h^2 + 5\left(y_1^2+y_1y_2+y_2^2\right)\right]\\ &-i\left(y_1+y_2\right)\left[7h^2\left(y_1^2+y_2^2\right) - 5\left(y_1^4+y_1^2y_2^2+y_2^4\right)\right]\\ &+d\Big[28h^2\left(y_1^2+y_1y_2+y_2^2\right) - 18\left(y_1^4+y_1^3y_2+y_1^2y_2^2+y_1y_2^3+y_2^4\right)\Big]\Big\}\end{aligned}$$

$$\begin{aligned}q_{55} = q_{00}\frac{1}{16}\sqrt{\frac{11}{7}}&\left[-2d + i\left(y_1+y_2\right)\right]\left[d^2 - y_1^2 + y_1y_2 - y_2^2 - id\left(y_1+y_2\right)\right]\\ &\times\left[3d^2 - y_1^2 - y_1y_2 - y_2^2 - 3id\left(y_1+y_2\right)\right]\end{aligned}$$

2.6. Trapezoidal slab

The trapezoidal slab has reflection symmetry about the $\hat{x}$-$\hat{y}$ plane and a thickness $t > 0$. Two of its sides are parallel to $\hat{y}$ with lengths w_1 and w_2. It is symmetric about $\hat{x}$, i.e., the two parallel sides are centered at $y = 0$ with separation $h > 0$. We define $w \equiv (w_1 + w_2)/2$ and $\Delta_w \equiv (w_1 - w_2)/w$.

$$q_{00} = \frac{\rho}{2\sqrt{\pi}}htw$$

$$q_{11} = q_{00}\frac{1}{4\sqrt{6}}h\left(\Delta_w - 6\right)$$

$$q_{20} = q_{00}\frac{\sqrt{5}}{96}\left[4\Delta_w h^2 - \Delta_w^2 w^2 - 4\left(4h^2 - 2t^2 + w^2\right)\right]$$

$$q_{22} = -q_{00}\frac{1}{32}\sqrt{\frac{5}{6}}\left[4\Delta_w h^2 + \Delta_w^2 w^2 + 4\left(-4h^2 + w^2\right)\right]$$

$$\begin{aligned} q_{31} = -q_{00}\frac{1}{3840}\sqrt{\frac{7}{3}}h\Big[&3\Delta_w^3 w^2 - 30\Delta_w^2 w^2 - 120\left(6h^2 - 4t^2 + w^2\right) \\ &+4\Delta_w\left(54h^2 - 20t^2 + 15w^2\right)\Big] \end{aligned}$$

$$q_{33} = q_{00}\frac{1}{256}\sqrt{\frac{7}{5}}h\Big[-\Delta_w^3 w^2 + 10\Delta_w^2 w^2 + 4\Delta_w\left(6h^2 - 5w^2\right) + 40\left(-2h^2 + w^2\right)\Big]$$

$$\begin{aligned} q_{40} = q_{00}\frac{1}{10240}\Big\{&-24\Delta_w^3 w^2 h^2 + 9\Delta_w^4 w^4 - 32\Delta_w h^2\left(24h^2 - 20t^2 + 15w^2\right) \\ &+8\Delta_w^2 w^2\left(24h^2 - 20t^2 + 15w^2\right) + 16\Big[144h^4 + 24t^4 - 40t^2 w^2 + 9w^4 \\ &-40h^2\left(4t^2 - w^2\right)\Big]\Big\} \end{aligned}$$

$$\begin{aligned} q_{42} = q_{00}\frac{1}{1024\sqrt{10}}\Big[&32\Delta_w\left(8h^4 - 5h^2 t^2\right) + 3\Delta_w^4 w^4 \\ &+40\Delta_w^2 w^2\left(-t^2 + w^2\right) + 16\left(-4h^2 + w^2\right)\left(12h^2 - 10t^2 + 3w^2\right)\Big] \end{aligned}$$

$$\begin{aligned} q_{44} = q_{00}\frac{1}{2048}\sqrt{\frac{7}{10}}\Big[&24\Delta_w^3 w^2 h^2 + 3\Delta_w^4 w^4 + \Delta_w\left(-256h^4 + 480h^2 w^2\right) \\ &+\Delta_w^2 w^2\left(-192h^2 + 40w^2\right) + 16\left(48h^4 - 40h^2 w^2 + 3w^4\right)\Big] \end{aligned}$$

$$\begin{aligned} q_{51} = q_{00}\frac{1}{57344}\sqrt{\frac{11}{30}}h\Big\{&3\Delta_w^5 w^4 - 42\Delta_w^4 w^4 + 8\Delta_w^3 w^2\Big(26h^2 - 14t^2 \\ &+21w^2\Big) - 224\Big[80h^4 + 24t^4 - 20t^2 w^2 + 3w^4 \\ &-20h^2\left(6t^2 - w^2\right)\Big] - 112\Delta_w^2 w^2\left[14h^2 + 5\left(-2t^2 + w^2\right)\right] \\ &+16\Delta_w\Big[400h^4 - 252h^2\left(2t^2 - w^2\right) + 7\left(8t^4 - 20t^2 w^2 + 5w^4\right)\Big]\Big\} \end{aligned}$$

$$\begin{aligned} q_{53} = -q_{00}\frac{1}{49152}\sqrt{\frac{11}{35}}h\Big\{&-9\Delta_w^5 w^4 + 126\Delta_w^4 w^4 + 112\Delta_w^2 w^2\Big(14h^2 \\ &-20t^2 + 15w^2\Big) - 8\Delta_w^3 w^2\left(26h^2 - 28t^2 + 63w^2\right) \\ &-224\left[80h^4 + 40t^2 w^2 - 9w^4 - 20h^2\left(4t^2 + w^2\right)\right] \\ &+16\Delta_w\left[400h^4 - 84h^2\left(4t^2 + 3w^2\right) + 35\left(8t^2 w^2 - 3w^4\right)\right]\Big\} \end{aligned}$$

$$\begin{aligned} q_{55} = q_{00}\frac{1}{16384}\sqrt{\frac{11}{7}}h\Big[&3\Delta_w^5 w^4 - 42\Delta_w^4 w^4 + \Delta_w^3 w^2\left(-208h^2 + 168w^2\right) \\ &+112\Delta_w^2 w^2\left(14h^2 - 5w^2\right) + 16\Delta_w\left(80h^4 - 252h^2 w^2 + 35w^4\right) \\ &-224\left(16h^4 - 20h^2 w^2 + 3w^4\right)\Big] \end{aligned}$$

2.7. Cylindrical section

The cylindrical section has a cross section consisting of the space between a chord of a circle (parallel to the $\hat{y}$ axis) and the arc the chord subtends. This object has reflection symmetry about the $\hat{x}$-$\hat{y}$ plane and is specified by its thickness, $t > 0$, the circle radius, r, and the distance along the $\hat{x}$ axis, $d > 0$, from the center of the circle to the chord. We define $s \equiv \sqrt{1 - d^2/r^2}$.

$$q_{00} = \frac{\rho}{2\sqrt{\pi}} tr \left[-Sd + r\cos^{-1}\left(\frac{d}{r}\right) \right]$$

$$q_{11} = \frac{\rho}{\sqrt{6\pi}} Str\left(d^2 - r^2\right)$$

$$q_{20} = \frac{\rho}{24}\sqrt{\frac{5}{\pi}} tr \left[Sd\left(2d^2 - t^2 + r^2\right) + r\left(t^2 - 3r^2\right)\cos^{-1}\left(\frac{d}{r}\right) \right]$$

$$q_{22} = \frac{\rho}{2}\sqrt{\frac{5}{6\pi}} Sdtr\left(r^2 - d^2\right)$$

$$q_{31} = \frac{\rho}{60}\sqrt{\frac{7}{3\pi}} Str\left(r^2 - d^2\right)\left(6d^2 - 5h^2 + 9r^2\right)$$

$$q_{33} = \frac{\rho}{12} tr\sqrt{\frac{7}{5\pi}} S\left(6d^4 - 7d^2r^2 + r^4\right)$$

$$q_{40} = \frac{\rho}{160\sqrt{\pi}} tr \Bigg\{ - \Bigg[Sd\big(16d^4 - 20d^2t^2 + 3t^4 + 2\left(4d^2 - 5t^2\right)r^2 + 6r^4\big) \Bigg] + 3r\left(t^4 - 10t^2r^2 + 10r^4\right)\cos^{-1}\left(\frac{d}{r}\right) \Bigg\}$$

$$q_{42} = \frac{\rho}{8\sqrt{10\pi}} Sdtr\left(d^2 - r^2\right)\left(4d^2 - 5t^2 + 6r^2\right)$$

$$q_{44} = -\frac{\rho}{8} dtr\sqrt{\frac{7}{10\pi}} S\left(8d^4 - 11d^2r^2 + 3r^4\right)$$

$$q_{51} = \frac{\rho}{112} tr\left(d^2 - r^2\right)\sqrt{\frac{11}{30\pi}} S\Big[16d^4 - 28d^2t^2 + 7t^4 + 6\left(4d^2 - 7t^2\right)r^2 + 30r^4\Big]$$

$$q_{53} = \frac{\rho}{144} tr\left(d^2 - r^2\right)\sqrt{\frac{11}{35\pi}} S\Big[- 48d^4 - 14t^2r^2 + 15r^4 + 12d^2\left(7t^2 - 6r^2\right)\Big]$$

$$q_{55} = \frac{\rho}{80} tr\left(d^2 - r^2\right)\sqrt{\frac{11}{7\pi}} S\left(80d^4 - 48d^2r^2 + 3r^4\right)$$

2.8. Right tetrahedron

This shape consists of a tetrahedron having three mutually perpendicular triangular faces that meet at the origin. The fourth triangular face is defined by points at coordinates x, y and z along the $\hat{x}$, $\hat{y}$, and $\hat{z}$ axes, respectively.

$$q_{00} = \frac{\rho xyz}{12\sqrt{\pi}}$$

$$q_{10} = q_{00}\frac{\sqrt{3}}{4} z$$

$$q_{11} = -q_{00}\frac{3}{4\sqrt{6}}\left(x - iy\right)$$

$$q_{20} = -q_{00}\frac{1}{4\sqrt{5}}\left(x^2 + y^2 - 2z^2\right)$$

$$q_{21} = -q_{00}\frac{3}{4\sqrt{30}} z\left(x - iy\right)$$

$$q_{22} = q_{00}\frac{3}{4\sqrt{30}}\left(x^2 - ixy - y^2\right)$$

$$q_{30} = -q_{00}\frac{\sqrt{7}}{40} z\left(x^2 + y^2 - 2z^2\right)$$

$$q_{31} = q_{00}\frac{1}{80}\sqrt{\frac{7}{3}}\left[3x^3 - ix^2y + xy^2 - 3iy^3 - 4\left(x - iy\right)z^2\right]$$

$$q_{32} = q_{00}\frac{1}{8}\sqrt{\frac{7}{30}} z\left(x^2 - ixy - y^2\right)$$

$$q_{33} = -q_{00}\frac{1}{16}\sqrt{\frac{7}{5}}\left(x^2 - y^2\right)\left(x - iy\right)$$

$$q_{40} = q_{00}\frac{3}{280}\left[3x^4 + x^2y^2 + 3y^4 - 4\left(x^2 + y^2\right)z^2 + 8z^4\right]$$

$$q_{41} = q_{00}\frac{3}{112\sqrt{5}} z\left[3x^3 - ix^2y - 3iy^3 + 4iyz^2 + x\left(y^2 - 4z^2\right)\right]$$

$$q_{42} = -q_{00}\frac{3}{56\sqrt{10}}\left[2x^4 - ix^3y - ixy^3 - 2y^4 - 2\left(x^2 - ixy - y^2\right)z^2\right]$$

$$q_{43} = -q_{00}\frac{3}{16\sqrt{35}} z\left(x^2 - y^2\right)\left(x - iy\right)$$

$$q_{44} = q_{00}\frac{3}{8\sqrt{70}}\left(x^4 - ix^3y - x^2y^2 + ixy^3 + y^4\right)$$

$$q_{50} = q_{00}\frac{1}{4486}\sqrt{11}\, z\left[3x^4 + x^2y^2 + 3y^4 - 4\left(x^2 + y^2\right)z^2 + 8z^4\right]$$

$$\begin{aligned} q_{51} = -q_{00}\frac{3}{448}\sqrt{\frac{11}{30}}\Big[&5x^5 - ix^4y + x^3y^2 - ix^2y^3 + xy^4 - 5iy^5 - \\ &2\left(x + iy\right)\left(3x^2 - 4ixy - 3y^2\right)z^2 + 8\left(x - iy\right)z^4\Big] \end{aligned}$$

$$q_{52} = q_{00}\frac{3}{64}\sqrt{\frac{11}{210}} z\left[-2x^4 + ix^3y + ixy^3 + 2y^4 + 2\left(x^2 - ixy - y^2\right)z^2\right]$$

$$q_{53} = q_{00}\frac{1}{128}\sqrt{\frac{11}{35}}\left[5x^5 - 3ix^4y - x^3y^2 - ix^2y^3 - 3xy^4 + 5iy^5 - 4\left(x^2 - y^2\right)\left(x - iy\right)z^2\right]$$

$$q_{54} = q_{00}\frac{3}{64}\sqrt{\frac{11}{70}} z\left(x^4 - ix^3y - x^2y^2 + ixy^3 + y^4\right)z^2$$

$$q_{55} = -q_{00}\frac{3}{128}\sqrt{\frac{11}{7}}\left(x - iy\right)\left[\left(x^2 - y^2\right)^2 + x^2y^2\right]$$

2.9. Cylinder hole

This shape consists of the volume that would be removed by drilling a hole of radius r into a cylinder of radius R. The symmetry axis of the hole is along $\hat{z}$, and the cylinder has its symmetry axis along $\hat{y}$. The moments require the hypergeometric function ${}_2F_1(a, b; c; x)$.

$$q_{00} = \rho\sqrt{\pi} r^2 R\, {}_2F_1\left(-\frac{1}{2}, \frac{1}{2}; 2; \frac{r^2}{R^2}\right)$$

$$q_{20} = -\frac{\rho}{6}\sqrt{5\pi}r^2R\Big[\left(r^2-2R^2\right)\ {}_2F_1\left(-\frac{1}{2},\frac{1}{2};2;\frac{r^2}{R^2}\right)+r^2\ {}_2F_1\left(-\frac{1}{2},\frac{3}{2};3;\frac{r^2}{R^2}\right)\Big]$$

$$q_{22} = \rho\sqrt{\frac{15\pi}{2}}r^2R\ {}_2F_1\left(-\frac{1}{2},\frac{1}{2};2;\frac{r^2}{R^2}\right)$$

$$\begin{aligned} q_{40} = \frac{\rho}{40}\sqrt{\pi}rR\Big[&\left(9r^5-40r^3R^2+24rR^4\right)\ {}_2F_1\left(-\frac{1}{2},\frac{1}{2};2;\frac{r^2}{R^2}\right) \\ &+\left(13r^5-32r^3R^2\right)\ {}_2F_1\left(-\frac{1}{2},\frac{3}{2};3;\frac{r^2}{R^2}\right)+16r^5\ {}_2F_1\left(-\frac{1}{2},\frac{5}{2};4;\frac{r^2}{R^2}\right)\Big]\end{aligned}$$

$$\begin{aligned} q_{42} = \frac{\rho}{8}\sqrt{\frac{\pi}{10}}r^4R\Big[&\left(6r^2-20R^2\right)\ {}_2F_1\left(-\frac{1}{2},\frac{1}{2};2;\frac{r^2}{R^2}\right) \\ &+2\left(r^2+10R^2\right)\ {}_2F_1\left(-\frac{1}{2},\frac{3}{2};3;\frac{r^2}{R^2}\right)-13r^2\ {}_2F_1\left(-\frac{1}{2},\frac{5}{2};4;\frac{r^2}{R^2}\right)\Big]\end{aligned}$$

$$q_{44} = \frac{3\rho}{2}\sqrt{\frac{35\pi}{2}}r^2R\ {}_2F_1\left(-\frac{1}{2},\frac{1}{2};2;\frac{r^2}{R^2}\right)$$

2.10. Plate hole

This shape consists of the volume that would be removed by drilling a hole of radius r through a parallel-sided plate of thickness t. The plate is centered on the $\hat{x}$-$\hat{y}$ plane. The hole axis, which passes through the origin, lies in the $\hat{x}$-$\hat{z}$ plane at an angle θ measured from $\hat{z}$, where $-\pi/2 < \theta < \pi/2$. We define $s \equiv \tan\theta$.

$$q_{00} = \frac{\rho t r^2\sqrt{\pi}(1+s^2)}{2}$$

$$q_{20} = q_{00}\frac{\sqrt{5}}{24}\left[3r^2(s^2+2)+t^2(s^2-2)\right]$$

$$q_{21} = -\ q_{00}\frac{1}{4}\sqrt{\frac{5}{6}}t^2s$$

$$q_{22} = -\ q_{00}\frac{1}{8}\sqrt{\frac{5}{6}}s^2(3r^2+t^2)$$

$$q_{40} = -\ q_{00}\frac{3}{640}\left[10r^4(8+8s^2+3s^4)+10t^2r^2(3s^4-8)+t^4\left(8-24s^2+3s^4\right)\right]$$

$$q_{41} = -\ q_{00}\frac{3}{64\sqrt{5}}t^2s\left[5r^2\left(4+3s^2\right)+t^2\left(3s^2-4\right)\right]$$

$$q_{42} = q_{00}\frac{3}{64}\sqrt{\frac{1}{10}}s^2\left[10r^4\left(s^2+2\right)+10r^2s^2t^2+t^4\left(s^2-6\right)\right]$$

$$q_{43} = q_{00}\frac{3}{64}\sqrt{\frac{7}{5}}s^3t^2\left(5r^2+t^2\right)$$

$$q_{44} = -\ q_{00}\frac{3}{128}\sqrt{\frac{7}{10}}s^4\left(10r^4+10r^2t^2+t^4\right)\ ,$$

2.11. Pyramid

This shape consists of a symmetric pyramid whose base with side lengths x and y lies in the $\hat{x}$-$\hat{y}$ plane centered about the origin, and whose apex is at $(x, y, z) = (0, 0, h)$.

$$q_{00} = \frac{\rho h x y}{6\sqrt{\pi}}$$

$$q_{10} = q_{00} \frac{\sqrt{3} h}{4}$$

$$q_{20} = q_{00} \frac{4h^2 - x^2 - y^2}{8\sqrt{5}}$$

$$q_{22} = q_{00} \sqrt{\frac{3}{10}} \frac{x^2 - y^2}{8}$$

$$q_{30} = q_{00} \frac{\sqrt{7}}{80} h(4h^2 - x^2 - y^2)$$

$$q_{32} = q_{00} \sqrt{\frac{7}{30}} \frac{h(x^2 - y^2)}{16}$$

$$q_{40} = q_{00} \frac{3}{4480} \left[128h^4 + 9x^4 + 10x^2y^2 + 9y^4 - 32h^2(x^2 + y^2)\right]$$

$$q_{42} = q_{00} \frac{3}{448\sqrt{10}} (x^2 - y^2) \left[8h^2 - 3(x^2 + y^2)\right]$$

$$q_{44} = q_{00} \frac{3}{128\sqrt{70}} \left[3x^4 - 10x^2y^2 + 3y^2\right]$$

$$q_{50} = q_{00} \frac{\sqrt{11}}{7168} h \left[128h^4 + (x^4 + 10x^2y^2 + 9y^2 - 32h^2(x^2 + y^2)\right]$$

$$q_{52} = q_{00} \frac{1}{512} \sqrt{\frac{33}{70}} h(x^2 - y^2) \left[8h^2 - 3(x^2 + y^2)\right]$$

$$q_{54} = q_{00} \frac{3}{1024} \sqrt{\frac{11}{70}} \left(3x^4 - 10x^2y^2 + 3y^4\right) \ ,$$

3. Gravitational force between two rectangular solids at arbitrary separations

In situations where the test masses are in close proximity, the multipole formulation is not valid. In such cases one can use a rectangular mesh to model complex objects. We present here an analytic expression for the force between two parallel-sided rectangular solids at arbitrary separations. Mass 2 has density ρ_2 and is centered at the origin with side half-lengths (a_2, b_2, c_2). Mass 1 has density ρ_2 and is centered at $(\Delta x, \Delta y, \Delta z)$ with side half-lengths (a_1, b_1, c_1).

$$F_z = -G\rho_1\rho_2 \left[I_1^2 - I_0^2 - I_3^2 + I_2^2 - y_0 \left(I_1^1 - I_0^1\right) + (y_1 - y_0)(I_2^1 - I_1^1) + y_3(I_3^1 - I_2^1)\right] , \quad (1)$$

where $I_{\rm n}^1$, $I_{\rm n}^2$, and the $y_{\rm i}$ are derived in the Appendix. Similar expressions yield F_x and F_y. Although the complete expressions are quite long they can be evaluated very quickly on modern computers. The precision attainable with our expressions may be very useful when analyzing the gravitational balancing of drag-free spacecraft, such as those envisioned for LISA[4]. Our expression may also find use in accurately modeling the Newtonian forces between rectangular test masses in future short-range tests of gravity.

Appendix: derivation of the force between two rectangular solids

The z-component of the gravitational force on mass 2 is given by the 6-dimensional integral

$$F_z = -G\rho_1\rho_2 \int_{\Delta x-a_1}^{\Delta x+a_1}\int_{\Delta y-b_1}^{\Delta y+b_1}\int_{\Delta z-c_1}^{\Delta z+c_1}\int_{-a_2}^{a_2}\int_{-b_2}^{b_2}\int_{-c_2}^{c_2} \frac{\mathrm{d}U}{\mathrm{d}(z_1-z_2)} \mathrm{d}z_2\,\mathrm{d}y_2\,\mathrm{d}x_2\,\mathrm{d}z_1\,\mathrm{d}y_1\,\mathrm{d}x_1\ ,$$

where $U = \left[(x_1-x_2)^2+(y_1-y_2)^2+(z_1-z_2)^2\right]^{-1/2}$. We transform the coordinates into sums and differences. For example,

$$\begin{aligned} z &= z_1 - z_2 \\ z_s &= z_1 + z_2 \\ \mathrm{d}z\,\mathrm{d}z_s &= 2\,\mathrm{d}z_1\,\mathrm{d}z_2\ . \end{aligned}$$

Integration over z_s is trivial because the function is independent of this variable. The z-integrations give

$$\begin{aligned} \int \mathrm{d}z_1 \int \mathrm{d}z_2 \frac{\mathrm{d}U}{\mathrm{d}(z_1-z_2)} &= \frac{1}{2}\int \mathrm{d}z \int \mathrm{d}z_s \frac{\mathrm{d}U}{\mathrm{d}z} = \int_{z_2}^{z_3} U\mathrm{d}z - \int_{z_0}^{z_1} U\mathrm{d}z \\ &= F_0(x,y) + F_3(x,y) - F_1(x,y) - F_2(x,y)\ , \end{aligned}$$

where

$$F_{\rm i}(x,y) \equiv \ln\left(z_{\rm i} + \sqrt{x^2+y^2+z_{\rm i}^2}\right)\ .$$

The limits are given by

$$\begin{aligned} z_0 &= \Delta z - c_1 - c_2 \\ z_1 &= \mathrm{Min}\,(\Delta \mathrm{z} + \mathrm{c}_1 - \mathrm{c}_2,\ \Delta \mathrm{z} - \mathrm{c}_1 + \mathrm{c}_2) \\ z_2 &= \mathrm{Max}\,(\Delta \mathrm{z} + \mathrm{c}_1 - \mathrm{c}_2,\ \Delta \mathrm{z} - \mathrm{c}_1 + \mathrm{c}_2) \\ z_3 &= \Delta z + c_1 + c_2\ . \end{aligned}$$

Analogous expressions define the limits for the other coordinates. The x integrand involves terms such as $F_{\rm i}(x,y)$ and $x\,F_{\rm i}(x,y)$, which lead to

$$\begin{aligned} G^1_{\rm ij}(y) \equiv \int^{x_{\rm j}} \mathrm{d}x\, F_{\rm i}(x,y) = x_{\rm j}\left[\ln\left(z_{\rm i}+r_{\rm ij}\right)-1\right] + y\left[\tan^{-1}\!\left(\frac{x_{\rm j}}{y}\right) - \tan^{-1}\!\left(\frac{x_{\rm j} z_{\rm i}}{y\, r_{\rm ij}}\right)\right] \\ + z_{\rm i}\ln\left(x_{\rm j}+r_{\rm ij}\right) \end{aligned}$$

and

$$G^2_{\rm ij}(y) \equiv \int^{x_{\rm j}} \mathrm{d}x\, x\, F_{\rm i}(x,y) = \frac{1}{4}\left[2\, z_{\rm i}\, r_{\rm ij} - r_{\rm ij}^2 + 2\,\rho_{\rm j}^2\, \ln\left(z_{\rm i}+r_{\rm ij}\right)\right],$$

where $r_{\rm ij} = \sqrt{x_{\rm j}^2+y^2+z_{\rm i}^2}$ and $\rho_{\rm j} = \sqrt{x_{\rm j}^2+y^2}$. After performing the trivial integrals over x_s and y_s, five of the six integrations are complete. The final step is the integration over y, which contains the terms

$$\begin{aligned} \int^{y} G^1_{\rm ij}(y')\,\mathrm{d}y' = x_{\rm j}\, y\left[\ln\left(z_{\rm i}+r_{\rm ij}\right) - \frac{3}{2}\right] + y\, z_{\rm i}\left[\ln\left(x_{\rm j}+r_{\rm ij}\right)-1\right] \\ + \frac{y^2}{2}\left[\tan^{-1}\!\left(\frac{x_{\rm j}}{y}\right) - \tan^{-1}\!\left(\frac{x_{\rm j}\, z_{\rm i}}{y\, r_{\rm ij}}\right)\right] + \frac{z_{\rm i}^2}{2}\left[2\,\tan^{-1}\!\left(\frac{y}{z_{\rm i}}\right)\right. \end{aligned}$$

$$- \tan^{-1}\!\left(\frac{x_\mathrm{j}\, y}{z_\mathrm{i}\, r_\mathrm{ij}}\right)\Bigg] + \frac{x_\mathrm{j}^2}{2}\left[\tan^{-1}\!\left(\frac{y}{x_\mathrm{j}}\right) - \tan^{-1}\!\left(\frac{y\, z_\mathrm{i}}{x_\mathrm{j}\, r_\mathrm{ij}}\right)\right]$$

$$+ x_\mathrm{j}\, z_\mathrm{i}\, \ln\left(y + r_\mathrm{ij}\right)\ ,$$

$$\int^y y'\, G^1_\mathrm{ij}(y')\, \mathrm{d}y' = \frac{1}{12}\Big\{-3\, r_\mathrm{ij}^2\, (x_\mathrm{j} + z_\mathrm{i}) - 4\, x_\mathrm{j}(y^2 - z_\mathrm{i}\, r_\mathrm{ij}) + 6\, y^2\, z_\mathrm{i}\, \ln(x_\mathrm{j} + r_\mathrm{ij})$$

$$+4\, y^3\left[\tan^{-1}\!\left(\frac{x_\mathrm{j}}{y}\right) - \tan^{-1}\!\left(\frac{x_\mathrm{j}\, z_\mathrm{i}}{y\, r_\mathrm{ij}}\right)\right] +6\, x_\mathrm{j}\, y^2\, \ln(z_\mathrm{i} + r_\mathrm{ij})$$

$$-2\, z_\mathrm{i}^3\, \ln\left(\frac{36\,(x_\mathrm{j}^2 + z_\mathrm{i}^2)}{x_\mathrm{j}^4\, z_\mathrm{i}^6\,(y^2 + z_\mathrm{i}^2)(x_\mathrm{j} + r_\mathrm{ij})}\right) -2\, x_\mathrm{j}^3\, \ln\left(\frac{36(x_\mathrm{j}^2 + z_\mathrm{i}^2)}{x_\mathrm{j}^6\, z_\mathrm{i}^4 (r_\mathrm{ij} + z_\mathrm{i})}\right)\Bigg\}\ ,$$

$$\int^y G^2_\mathrm{ij}(y')\, \mathrm{d}y' = \frac{1}{36}\Bigg(6\,\Bigg\{2\, x_\mathrm{j}^3\left[\tan^{-1}\!\left(\frac{y}{x_\mathrm{j}}\right) - \tan^{-1}\!\left(\frac{y\, z_\mathrm{i}}{x_\mathrm{j}\, r_\mathrm{ij}}\right)\right] + z_\mathrm{i}\,(3x_\mathrm{j}^2 + z_\mathrm{i}^2)\ln(y + r_\mathrm{ij})$$

$$+y\,(3x_\mathrm{j}^2 + y^2)\ln(z_\mathrm{i} + r_\mathrm{ij})\Bigg\} - y\,(5r_\mathrm{ij} + 16x_\mathrm{j}^2 + 4z_\mathrm{i}^2 - 12r_\mathrm{ij}z_\mathrm{i})\Bigg)\ ,$$

and

$$\int^y y'\, G^2_\mathrm{ij}(y')\, \mathrm{d}y' = \frac{1}{96}\left[8z_\mathrm{i}^3\, r_\mathrm{ij} - \left(3\rho_\mathrm{j}^2 + z_\mathrm{i}^2\right)^2 - 6\rho_\mathrm{j}^2 z_\mathrm{i}^2 + 12\rho_\mathrm{j}^4 \ln(z_\mathrm{i} + r_\mathrm{ij})\right]\ .$$

Finally, we define

$$H_\mathrm{k}(y) \equiv G^2_\mathrm{k1}(y) - G^2_\mathrm{k0}(y) - x_0\,(G^1_\mathrm{k1}(y) - G^1_\mathrm{k0}(y)) + (x_1 - x_0)\,(G^1_\mathrm{k2}(y) - G^1_\mathrm{k1}(y))$$

$$+x_3(G^1_\mathrm{k3}(y) - G^1_\mathrm{k2}(y)) - G^2_\mathrm{k3}(y) + G^2_\mathrm{k2}(y)$$

and

$$I^1_\mathrm{n} \equiv \int^{y_\mathrm{n}} \mathrm{d}y\,[H_0(y) + H_3(y) - H_1(y) - H_2(y)]$$

$$I^2_\mathrm{n} \equiv \int^{y_\mathrm{n}} y\, \mathrm{d}y\,[H_0(y) + H_3(y) - H_1(y) - H_2(y)]\ ,$$

which lead to Equation 1.

Acknowledgments

This work was supported in part by National Science Foundation grants PHY-9602494 and PHY-0355012. CDH would like to thank M. Armano for carefully checking the rectangular solid force calculation.

References

[1] Y. Su et al., Phys, Rev. D **50**, 3614 (1994).
[2] G.L. Smith et al., Phys. Rev. D **61**, 022001-1 (2000).
[3] C. D'Urso and E.G. Adelberger, Phys. Rev. D **55**, 7970 (1997).
[4] M. Armano et al., Class. Quantum Grav. **22** S501 (2005).